

Test of a simple and flexible S_8 model molecule in α - S_8 crystals

C. Pastorino and Z. Gamba

Department of Physics, Comisión Nacional de Energía Atómica,

Av. Libertador 8250, (1429) Buenos Aires, Argentina. e-mails:

clopasto@cnea.gov.ar and gamba@cnea.gov.ar

α - S_8 is the most stable crystalline form, at ambient pressure and temperature (STP), of elemental sulfur. In this paper we analyze the zero pressure low temperature part of the phase diagram of this crystal, in order to test a simple and flexible model molecule. The calculations consist in a series of molecular dynamics (MD) simulations, performed in the constant pressure-constant temperature ensemble. Our calculations show that this model, that gives good results for three crystalline phases at STP and $T \gtrsim 300\text{K}$, fails at low temperatures, predicting a structural phase transition at 200K where there should be none.

Introduction:

Elemental sulfur is of relevance in geophysics, astrophysics, material sciences and massively used in industry [1,2]. Recently, a high pressure superconductor transition, with $T_c=17\text{K}$, has been found [3].

Elemental sulfur is widely known by the unusual characteristics of its liquid phase. Numerous works have been dedicated to the study of its liquid-liquid (λ) phase transition, due to a large variation of its molecular composition. The λ transition is of relevance in numerous fields, volcanology is an example [4–6]. The liquid phase is also of enormous theoretical interest, with studies ranging from models to describe the λ transition [7–9], to molecular dynamics (MD) simulations of the liquid [10–12] and to calculations of the stability and structure of isolated S_n molecules [13–16].

In spite of the numerous experimental and theoretical studies on the liquid phase, the

phase diagram and properties of the crystalline phases of elemental sulfur are far from well known [1,2]. A simple inter- and intramolecular potential, able to reproduce the structural and dynamical properties of the crystalline phases, should be very useful to understand the complex experimental data of sulfur crystals.

The most stable sulfur allotrope at ambient temperature and pressure (STP) is S_8 , a cyclic crown-shaped molecule, stable in solid, liquid and gas phases. Three crystalline phases have been clearly identified at zero pressure and $T \lesssim 400\text{K}$, being orthorhombic α - S_8 the most stable crystalline form at STP. Fig. 1a shows its "crankshaft" structure with four molecules in the primitive unit cell. The space group is D_{2h}^{24} (Fddd), with 16 molecules in this non-primitive orthorhombic cell. If α - S_8 is slowly heated, it shows a phase transition to monoclinic β - S_8 at 369K, which melts at 393K. [1]. Nevertheless, most α - S_8 crystals do not convert to β - S_8 [17], they melt, instead, at 385.8K [1,2].

Experimentally, the samples of pure α - S_8 can be easily identified because these crystals are greenish-yellow at 293K and turn white below 193K. Commercial sulfur, instead, contains small amounts of S_7 and conserves the yellow color at low temperatures [1]. The change of color is assumed to be due to a shift of the electronic band gap between valence and conduction bands [1,2].

The firsts molecular dynamics (MD) simulations of α - S_8 crystals were performed in the constant volume- constant energy ensemble, at normal pressure and at several temperatures [18], with the only aim to study the Raman and ir. intramolecular frequencies in a crystalline environment. Numerous works [19] have also been dedicated to the measurement and/or calculation, by lattice sums and in different approximations, of the $\mathbf{k}=0$ vibrational modes.

The first attempt to reproduce the zero pressure phase diagram of α -, β - and γ - S_8 crystals has been recently performed [20], using a simple and flexible model molecule, based in that proposed by Venutti et al. [18]. The model molecule reproduces the STP structures the three crystals, the measured heat of sublimation of α - S_8 , and the orientationally disordered high temperature plastic phase of β - S_8 crystals.

In this paper we study the zero pressure and low temperature phase diagram of α - S_8 crys-

tals, as given by the simple and flexible model molecule of ref. [18]. The calculations consist of a series of constant pressure - constant temperature classical MD simulations, at several temperatures and zero pressure. This MD algorithm is useful to locate structural phase transitions, or in the search of the most stable structure for a given set of thermodynamic parameters P and T.

Unfortunately, the model molecule of ref. [20] fails to reproduce the measured low temperature structure of α -S₈ crystals: a structural phase transition is found at 200K, where there should be none. Here we give a detailed account of the calculations performed at low temperatures on α -S₈ crystals, of a disordered sample, and of the reasons for the failure of the proposed model.

The inter- and intramolecular potential model:

The main problem found in this type of calculations is the inter- and intramolecular potential model of this flexible molecule. The model of our simulations is a slight modification of that proposed in the constant volume MD simulations of α -S₈ crystals, that successfully reproduced the available experimental data on the crystalline intramolecular frequencies [18]. The intermolecular potential model of ref. [18] was of the Lennard-Jones (LJ) atom-atom type. In our constant pressure- constant temperature MD calculations, these LJ parameters [18] were slightly changed to improve the fit of the calculated STP crystalline structure and configurational energy of α -S₈ to the experimental data. The parameters of our LJ potential model, for non-bonded S-S interactions are: $\varepsilon = 1.70 \text{ kJ/mol}$ and $\sigma = 3.39 \text{ \AA}$. The cut-off radius of atom-atom interactions is $12 \text{ \AA}$ and correction terms to the configurational energy and pressure, due to this finite value, are taken into account in the usual way: by integrating the contribution of a uniform distribution of atoms.

A similar procedure is followed for the intramolecular potentials of the bending and torsional angles, that are based on those of ref. [18]. Our flexible model molecule takes into account all the low frequency intramolecular modes that mix with lattice modes and can therefore be relevant to the onset of structural changes as a function of T and P values.

Since the stretching modes (from 400 to 500cm⁻¹) are always well above in energies than the rest of the vibrational modes (< 250cm⁻¹) and of the lattice modes (between 10 and ~100cm⁻¹), the S-S bond length of our model molecule is held constant at 2.0601Å. The bending and torsional intramolecular motions (from 150 to 250cm⁻¹), instead, are taken into account and their mix with lattice modes is allowed in a straightforward way.

The intramolecular potential for bending angles S-S-S is taken harmonic

$$V(\beta) = \frac{1}{2}C_{\beta}(\beta - \beta_0)^2,$$

with a force constant of $C_{\beta}=25700\text{Kk}_B/\text{rad}^2$ and $\beta_0=108\text{deg}$. The intramolecular potential for torsion (τ) angles is a double well:

$$V(\tau) = A_{\tau} + B_{\tau} \cos(\tau) + C_{\tau} \cos^2(\tau) + D_{\tau} \cos^3(\tau),$$

with $A_{\tau}=57.19\text{ k}_B$, $B_{\tau}=738.41\text{ k}_B$, $C_{\tau}=2297.90\text{ k}_B$ and $D_{\tau}=557.25\text{ k}_B$. These parameters describe a double well with minima at $\tau=\pm 98.8\text{deg}$., and a height barrier of about 12kJ/mol.

Calculations:

The structural and dynamical properties of α -S₈, are studied in the (N,P,T) ensemble, *via* a series of classical constant pressure - constant temperature MD simulations. The standard MD algorithm allows volume and shape fluctuations of the MD sample in order to balance the applied isotropic external pressure with the internal stresses [21], the temperature control of the sample follows the approach of Nosé [22,23]. The equations of motion of these flexible molecules are integrated using the Verlet algorithm for the atomic displacements and the Shake algorithm for the constant bond lengths constraints on each molecule [24,25]. The final MD algorithm is identical to that used in a study of black Newton films [26].

The starting point of our simulations is the experimental structure of α -S₈ at STP [27]. The sample was first equilibrated with a constant volume algorithm on a trajectory of 30000 time steps (of 0.01ps.). Afterwards, the runs in the (N,P,T) ensemble were performed by

decreasing and increasing the temperature (up to 450K) in steps of 25 or 50K. At each point of the phase diagram the sample is equilibrated for 20000 to 30000 time steps and measured in the following 10ps. Near the phase transitions the equilibration times were increased several times.

The sample of α -S₈ crystals consisted of 3x3x2 orthorhombic cells (288 molecules). Some points of the phase diagram were recalculated with a sample of 4x4x2 cells (512 molecules).

Results:

a) The α -phase:

α -S₈ crystallizes, at STP, in the orthorhombic space group F_d (D_{2h}^{24}) with 16 molecules in a face-centered cell, the lattice parameters at 300K are $a=10.4646\text{\AA}$, $b=12.8660\text{\AA}$ and $c=24.4860\text{\AA}$ [27]. The primitive cell contains 4 molecules at C_2 sites. The molecular rings lie parallel to the crystallographic c axis and are alternately oriented parallel to (110) and ($\bar{1}\bar{1}0$) crystalline planes. Fig.1a includes part of our sample at 300K and shows the structure in the [110] direction. At 300K our calculated lattice parameters are $a=10.34(10)\text{\AA}$, $b=13.20(3)\text{\AA}$ and $c=24.20(5)\text{\AA}$, the largest difference is found in the value of b axis, 2 % higher than its experimental value. The molecules location and orientation follow the experimental data [27]. At 300K the calculated value of configurational energy is -103.5(2)kJ/mol, from which the heat of sublimation can be estimated as 101.0kJ/mol and compares well with a measured value of 101.8(2)kJ/mol [28].

Figs. 2a and 2b show the calculated vibrational density of states of the isolated molecule and that of α -S₈ at 300K. They reproduce the experimental pattern, although the intramolecular modes are calculated about 10% higher than the experimental ones.

Figs. 3a and 3b show the calculated configurational energy and volume per molecule as a function of temperature, for both samples of α -S₈. Unfortunately, our orthorhombic α -S₈ sample of 288 molecules turns out to be unstable for $T \leq 200\text{K}$ and distorts to a monoclinic cell. The final molecular array closely resembles that of α -S₈ (Fig. 1b). Fig3c shows the

calculated angles of the unit cell as a function of temperature. This result was later on confirmed with a larger sample of 512 molecules (4x4x2 orthorhombic cells). Below 200K this large sample do not distort to monoclinic, but nevertheless its configurational energy and volume per molecule are higher than those for the sample of 288 molecules (Figs. 3a and 3b). The fluctuations of energy and volume, in the sample of 512 molecules, are also larger at $T \leq 200$ than those at 200K.

The coincidence of the experimental change of color at 193K and our calculated transition at 200K, made us review the experimental data on this phase. The usual reference quoted for the lattice parameters values, as a function of temperature, is ref. [30]. Their reported values were determined by X-ray powder diffractrometry, on different samples and by measuring only three lines at 8 temperatures between 100 and 300K. Our calculated powder diffraction patterns, not included here, are quite similar for the orthorhombic and monoclinic samples and no conclusive answer can be obtained from these data. Nevertheless, there is a careful single crystal determination of the structure of α -S₈ at 300 and 100K [31], where the crystalline charge distribution is measured. Both measured structures belong to the same crystalline space group and the molecular arrangement is identical. This study clearly disregard any possible structural phase transition and shows that the simple model molecule, in spite of the amount of experimental data that it is able to reproduce for $T \geq 200$ K, should still be improved [20].

b) Phase transitions and the liquid phase:

The experimental evidence shows that the phase transitions of S₈ crystals are obtained when defects are present, if they are not, metastable states can be maintained for days [1,2,17]. This evidence is valid for the $\alpha - \beta$ transition, and also for the solid-liquid phase transition [2]. In the last case, the disorder is generated because the S₈ molecules start to dissociate when the temperature is increased.

In our simulations, all calculated ordered samples of α -S₈ crystals melt at $T \geq 450$ K. That the disorder is essential to promote the solid-liquid phase transition can be checked by sim-

ulating a disordered sample of S_8 molecules. A cubic sample of 216 disordered S_8 molecules was studied in the range 350-450K, following the same procedure as for the other samples. In this case the MD algorithm allowed changes in the volume of the MD box, but not in its shape. Fig. 4 shows the calculated discontinuity in the configurational energy and volume per molecule at 400K, near that measured for α - S_8 (386K) and β - S_8 (393K) crystals. For $T \geq 400$ K the sample is liquid, as was determined by its diffusion constant. These simulations should be valid for temperatures above and near the transition, when the fraction of broken molecules is small, and show that the model molecule is useful in this range of temperatures.

The calculated distances between firsts neighbor atoms compares well with those determined from the pair distribution function $g_2(r)$ obtained by neutron measurements [29]. Fig. 4a includes our calculated bonded and non-bonded atom-atom distances of crystalline α - S_8 at 300K, Fig. 4b corresponds to our disordered 'liquid' sample at 425K. The 3th and 5th peaks at 3.9Å and 5.4Å, which are not found in the liquid phase, correspond to non-bonded distances. Their disappearance, observed in the liquid phase, cannot be unequivocally attributed to broken bonds [29].

Conclusions:

In this paper we study the zero pressure low temperature range of the phase diagram of S_8 crystals, as given by a simple and flexible model molecule. The molecule model, a simplified and slight modification of that proposed in ref. [18], is extremely simple and gives good account of many experimental facts for $T \geq 200$ K, including the experimental data on the structure and dynamics of α -, β - and γ - S_8 crystals around STP and the orientationally disordered plastic phase of β - S_8 crystals [20]. It also reproduces the solid-liquid phase transition, near the experimental temperature, but only for a disordered sample of S_8 molecules [20]. The last calculation is only of theoretical interest, and we do not intended, here, to simulate the real liquid phase of elemental sulfur. Nevertheless, the pair correlation function calculated in this theoretical 'liquid' shows that the 3th and 5th neighbours identified in solid

phases correspond to non-bonded atoms, and that their disappearance at high temperatures can be expected even if the molecules do not dissociate in short segments.

In spite of all the properties of S_8 crystals that the simple and flexible model molecule reproduces, below 200K this model is unable to calculate the experimental orthorhombic structure of α - S_8 . Although no structural phase transitions are measured for α - S_8 crystals, our sample of 288 molecules is unstable and transforms to a monoclinic structure.

Acknowledgement: The authors thank CONICET for the grant PIP 0859/98.

References:

- [1] R. Steudel, *Chemie unserer Zeit* **30** (1996) 226.
- [2] B. Meyer, *Chem. Rev.* **76** (1976) 367.
- [3] V. V. Struzhkri et al., *Nature* **390** (1997) 382.
- [4] C. Oppenheimer, *J. of Vulcanology and Geothermal Research*, **49** (1992) 1.
- [5] A. W. Hurst, H. M. Bibby, B. J. Scott and M. J. McGuinness, *J. of Vulcanology and Geothermal Research*, **46** (1991) 1.
- [6] B. Takano, H. Saitoh and E. Takano, *Geochem. J.* **28** (1994) 199.
- [7] A. V. Tobolsky and A. Eisenberg, *J. Am. Chem. Soc.* **82** (1960) 289.
- [8] J. C. Wheeler and P. Pfeuty, *Phys. Rev. A* **24**, (1981) 1050.
- [9] Y. Duda and M. F. Holovko, *Phys. Chem. Liq.* **35** (1997) 175.
- [10] F. H. Stillinger, T. H. Weber and R. A. LaViolette, *J. Chem. Phys.* **85** (1986) 6460.

- [11] F. H. Stillinger and T. H. Weber, J. Phys. Chem. **91** (1987) 4899.
- [12] J. S. Tse and D. D. Klug, Phys. Rev. **B 59** (1999) 34.
- [13] P. Bicaï, D. Changkui, X. Shangda and X. Chuanyun, Phys. Rev. **B 50** (1994) 17556.
- [14] A. S. Zyubin and S. A. Dembowsky, Solid State Comm. **89** (1994) 335.
- [15] D. Hohl, R. O. Jones, R. Car and M. Parrinello, J. Phys. Chem. **89** (1988) 6823.
- [16] M. Springborg and R. O. Jones, Phys. Rev. Letters **57** (1986) 1145.
- [17] D. E. Sands, J. Am. Chem. Soc. **87** (1965) 1396.
- [18] E. Venuti, G. Cardini and E. Castellucci, Chem. Phys. **165** (1992) 313.
- [19] M. Becucci, R. Bini, E. Castellucci, B. Eckert and H. J. Jodl, J. Phys. Chem. **101** (1997) 2132;
and references therein.
- [20] C. Pastorino and Z. Gamba, J. Chem. Phys., in press (1999); <http://xxx.lanl.gov/cond-mat/9910529>.
- [21] S. Nose and M. L. Klein, Mol. Phys. **50** (1983) 1055.
- [22] S. Nosé, Progress of Teor. Phys. Suppl. **103** (1991) 1.
- [23] J. P. Ryckaert and G. Ciccotti, J. Chem. Phys. **78**, (1983) 7368.
- [24] M. Ferrario and J. Ryckaert, Mol. Phys. **54** (1985) 587.
- [25] M. P. Allen and D. J. Tildesley, "Computer simulations of liquids", Clarendon Press, Oxford 1990.
- [26] Z. Gamba, J. Hautman, J. C. Shelley and M. L. Klein, Langmuir **8** (1992) 3155.
- [27] S. C. Abrahams, Acta Cryst. **8** (1955) 661.
- [28] G. B. Gothrie, P. W. Scott and G. Waddington, Acta Cryst. **76** (1953) 1488.

- [29] R. Winter, P. A. Egelstaff, W-C. Pilgrim and W. S. Howells, J. Phys.: Condens. Matter **2** (1990) SA215.
- [30] J. Wallis, I. Sigalas and S. Hart, J. Appl. Cryst. **19** (1986) 273.
- [31] P. Coppens, Y. W. Yang, R. H. Blessing, W. F. Cooper and F. K. Larsen, J. Am. Chem. Soc. **99** (1977) 760.

Figures:


Fig. 1: Calculated stable structures, as given by this model molecule: a) orthorhombic α -S₈ at 300K, b) monoclinic α' -S₈ at 150K.

Fig. 2: Vibrational density of states: a) Isolated molecule and b) α -S₈ crystal, both at 300K.

Fig. 3: Calculated (a) configurational energy and (b) volume per molecule of α -S₈ (large circles: sample of 288 molecules and small circles: sample of 512 molecules) and a disordered cubic sample (triangles: increasing T, squares: decreasing T) as a function of temperature. For $T \geq 400$ K, the disordered cubic sample is liquid. (c) Unit cell angles of the α -S₈ sample of 288 molecules. The lines are a guide to the eyes.

Fig. 4: Pair correlation functions: a) α -S₈ at 300K, b) our theoretical 'liquid' sample of S₈ molecules at 425K. Full lines: total atom-atom $g_2(r)$ functions, dotted lines: excluding intramolecular distances.

T=250K



T=150K

